

On the size of graphs without repeated cycle lengths[★]

Chunhui Lai^{a,1}

^a*School of Mathematics and Statistics,
Minnan Normal University, Zhangzhou, Fujian, P.R. China.*

Abstract

In 1975, P. Erdős proposed the problem of determining the maximum number $f(n)$ of edges in a graph with n vertices in which any two cycles are of different lengths. In this paper, it is proved that

$$f(n) \geq n + \frac{107}{3}t + \frac{7}{3}$$

for $t = 1260r + 169$ ($r \geq 1$) and $n \geq \frac{2119}{4}t^2 + 87978t + \frac{15957}{4}$. Consequently, $\liminf_{n \rightarrow \infty} \frac{f(n)-n}{\sqrt{n}} \geq \sqrt{2 + \frac{7654}{19071}}$, which is better than the previous bounds $\sqrt{2}$ [Y. Shi, Discrete Math. 71(1988), 57-71], $\sqrt{2.4}$ [C. Lai, Australas. J. Combin. 27(2003), 101-105]. The conjecture $\lim_{n \rightarrow \infty} \frac{f(n)-n}{\sqrt{n}} = \sqrt{2.4}$ is not true.

Key words: Graph, cycle, number of edges.

AMS 2000 MSC: 05C38, 05C35.

1 Introduction

Let $f(n)$ be the maximum number of edges in a graph on n vertices in which no two cycles have the same length. In 1975, Erdős raised the problem of

[★] Project supported by the National Science Foundation of China(Nos.11401290,61379021), NSF of Fujian(Nos.2015J01018, 2014J01020, 2016J01027, 2016J01673), Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering", Project of Fujian Education Department(Nos.JA15295, JJZ160455).

Email address: laich2011@msn.cn; laichunhui@mnnu.edu.cn (Chunhui Lai).

¹ Corresponding author

determining $f(n)$ (see Bondy and Murty [1], p.247, Problem 11). Shi [15] proved a lower bound.

Theorem 1 (Shi [15])

$$f(n) \geq n + [(\sqrt{8n - 23} + 1)/2]$$

for $n \geq 3$.

Chen, Lehel, Jacobson, and Shreve [3], Jia [5], Lai [6–8], Shi [16,18–20] obtained some additional related results.

Boros, Caro, Füredi and Yuster [2] proved an upper bound.

Theorem 2 (Boros, Caro, Füredi and Yuster [2]) For n sufficiently large,

$$f(n) < n + 1.98\sqrt{n}.$$

Lai [9] improved the lower bound.

Theorem 3 (Lai [9])

$$f(n) \geq n + \sqrt{2.4}\sqrt{n}(1 - o(1))$$

Lai [6,9] proposed the following conjectures:

Conjecture 4 (Lai [9])

$$\lim_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} = \sqrt{2.4}.$$

Conjecture 5 (Lai [6])

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \leq \sqrt{3}.$$

Markström [13] raised the following problem:

Problem 6 (Markström [13]) Determine the maximum number of edges in a hamiltonian graph on n vertices with no repeated cycle lengths.

Results for the maximum number of edges in a 2-connected graph on n vertices in which no two cycles have the same length can be found in [2,3,15].

Survey articles on this problem can be found in Tian [21], Zhang [24], Lai and Liu [10].

The progress of all 50 problems in [1] can be find in Locke [12].

A related topic concerns the Entringer problem, which is to determine which simple graphs have exactly one cycle of each length l , $3 \leq l \leq v$ (see problem 10 in [1]). This problem was posed in 1973 by R. C. Entringer. For developments on this topic, see [4,11,13,14,17,22,23].

In this paper, we construct a graph G having no two cycles with the same length which leads to the following result.

Theorem 7 Let $t = 1260r + 169$ ($r \geq 1$), then

$$f(n) \geq n + \frac{107}{3}t + \frac{7}{3}$$

for $n \geq \frac{2119}{4}t^2 + 87978t + \frac{15957}{4}$.

Hence conjecture 4 is not true.

2 Proof of theorem 7

Proof. Let $t = 1260r + 169$, $r \geq 1$, $n_t = \frac{2119}{4}t^2 + 87978t + \frac{15957}{4}$, $n \geq n_t$. We shall show that there exists a graph G on n vertices with $n + \frac{107}{3}t + \frac{7}{3}$ edges such that all cycles in G have distinct lengths.

Now we construct the graph G which consists of a number of subgraphs: B_i , $(0 \leq i \leq 20t, 27t \leq i \leq 28t + 64, 29t - 734 \leq i \leq 29t + 267, 30t - 531 \leq i \leq 30t + 57, 31t - 741 \leq i \leq 31t + 58, 32t - 740 \leq i \leq 32t + 57, 33t - 741 \leq i \leq 33t + 57, 34t - 741 \leq i \leq 34t + 52, 35t - 746 \leq i \leq 35t + 60, 36t - 738 \leq i \leq 36t + 60, 37t - 738 \leq i \leq 37t + 799, i = 20t + j(1 \leq j \leq \frac{t-7}{6}), i = 20t + \frac{t-1}{6} + j(1 \leq j \leq \frac{t-7}{6}), i = 21t + 2j + 1(0 \leq j \leq t-1), i = 21t + 2j(0 \leq j \leq \frac{t-1}{2}), i = 23t + 2j + 1(0 \leq j \leq \frac{t-1}{2}), \text{ and } i = 20t + \frac{t-1}{6}, i = 20t + \frac{t-1}{3} + \frac{t-1}{6} - 1, i = 20t + \frac{t-1}{3} + \frac{t-1}{6}, i = 20t + \frac{2t-2}{3}, i = 21t - 2, i = 21t - 1, i = 26t).$

Now we define these B_i s. These subgraphs all only have a common vertex x , otherwise their vertex sets are pairwise disjoint.

For $1 \leq i \leq \frac{t-7}{6}$, let the subgraph B_{20t+i} consists of a cycle

$$xa_i^1 a_i^2 \dots a_i^{\frac{62t-8}{3}+2i} x$$

and a path:

$$xa_{i,1}^1 a_{i,1}^2 \dots a_{i,1}^{\frac{59t-5}{6}} a_i^{\frac{61t-1}{6}+i}$$

Based the construction, B_{20t+i} contains exactly three cycles of lengths:

$$20t + i, 20t + \frac{t-1}{3} + i - 1, 20t + \frac{2t-2}{3} + 2i - 1.$$

For $1 \leq i \leq \frac{t-7}{6}$, let the subgraph $B_{20t+\frac{t-1}{6}+i}$ consists of a cycle

$$xb_i^1 b_i^2 \dots b_i^{\frac{62t-5}{3}+2i} x$$

and a path:

$$xb_{i,1}^1 b_{i,1}^2 \dots b_{i,1}^{10t-1} b_i^{\frac{61t-1}{6}+i}$$

Based the construction, $B_{20t+\frac{t-1}{6}+i}$ contains exactly three cycles of lengths:

$$20t + \frac{t-1}{6} + i, 20t + \frac{t-1}{3} + \frac{t-1}{6} + i, 20t + \frac{2t-2}{3} + 2i.$$

For $0 \leq i \leq t-1$, let the subgraph $B_{21t+2i+1}$ consists of a cycle

$$xu_i^1 u_i^2 \dots u_i^{25t+2i-1} x$$

and a path:

$$xu_{i,1}^1 u_{i,1}^2 \dots u_{i,1}^{(19t+2i-1)/2} u_i^{(23t+2i+1)/2}$$

Based the construction, $B_{21t+2i+1}$ contains exactly three cycles of lengths:

$$21t + 2i + 1, 23t + 2i, 25t + 2i.$$

For $0 \leq i \leq \frac{t-3}{2}$, let the subgraph B_{21t+2i} consists of a cycle

$$xv_i^1 v_i^2 \dots v_i^{25t+2i} x$$

and a path:

$$xv_{i,1}^1 v_{i,1}^2 \dots v_{i,1}^{9t+i-1} v_i^{12t+i}$$

Based the construction, B_{21t+2i} contains exactly three cycles of lengths:

$$21t + 2i, 22t + 2i + 1, 25t + 2i + 1.$$

For $i = \frac{t-1}{2}$, B_{21t+2i} is simply a cycle of length $22t - 1$.

For $0 \leq i \leq \frac{t-3}{2}$, let the subgraph $B_{23t+2i+1}$ consists of a cycle

$$xw_i^1w_i^2\dots w_i^{26t+2i+1}x$$

and a path:

$$xw_{i,1}^1w_{i,1}^2\dots w_{i,1}^{(21t+2i-1)/2}w_i^{(25t+2i+1)/2}$$

Based the construction, $B_{23t+2i+1}$ contains exactly three cycles of lengths:

$$23t + 2i + 1, 24t + 2i + 2, 26t + 2i + 2.$$

For $i = \frac{t-1}{2}$, $B_{23t+2i+1}$ is simply a cycle of length $24t$.

For $58 \leq i \leq t - 742$, let the subgraph $B_{27t+i-57}$ consists of a cycle

$$C_{27t+i-57} = xy_i^1y_i^2\dots y_i^{132t+11i+893}x$$

and ten paths sharing a common vertex x , the other end vertices are on the cycle $C_{27t+i-57}$:

$$\begin{aligned} & xy_{i,1}^1y_{i,1}^2\dots y_{i,1}^{(17t-1)/2}y_i^{(37t-115)/2+i} \\ & xy_{i,2}^1y_{i,2}^2\dots y_{i,2}^{(19t-1)/2}y_i^{(57t-103)/2+2i} \\ & xy_{i,3}^1y_{i,3}^2\dots y_{i,3}^{(19t-1)/2}y_i^{(77t+315)/2+3i} \\ & xy_{i,4}^1y_{i,4}^2\dots y_{i,4}^{(21t-1)/2}y_i^{(97t+313)/2+4i} \\ & xy_{i,5}^1y_{i,5}^2\dots y_{i,5}^{(21t-1)/2}y_i^{(117t+313)/2+5i} \\ & xy_{i,6}^1y_{i,6}^2\dots y_{i,6}^{(23t-1)/2}y_i^{(137t+311)/2+6i} \\ & xy_{i,7}^1y_{i,7}^2\dots y_{i,7}^{(23t-1)/2}y_i^{(157t+309)/2+7i} \\ & xy_{i,8}^1y_{i,8}^2\dots y_{i,8}^{(25t-1)/2}y_i^{(177t+297)/2+8i} \\ & xy_{i,9}^1y_{i,9}^2\dots y_{i,9}^{(25t-1)/2}y_i^{(197t+301)/2+9i} \\ & xy_{i,10}^1y_{i,10}^2\dots y_{i,10}^{(27t-1)/2}y_i^{(217t+305)/2+10i}. \end{aligned}$$

As a cycle with d chords contains $\binom{d+2}{2}$ distinct cycles, $B_{27t+i-57}$ contains exactly 66 cycles of lengths:

$$\begin{aligned}
& 27t + i - 57, & 28t + i + 7, & 29t + i + 210, & 30t + i, \\
& 31t + i + 1, & 32t + i, & 33t + i, & 34t + i - 5, \\
& 35t + i + 3, & 36t + i + 3, & 37t + i + 742, & 38t + 2i - 51, \\
& 38t + 2i + 216, & 40t + 2i + 209, & 40t + 2i, & 42t + 2i, \\
& 42t + 2i - 1, & 44t + 2i - 6, & 44t + 2i - 3, & 46t + 2i + 5, \\
& 46t + 2i + 744, & 48t + 3i + 158, & 49t + 3i + 215, & 50t + 3i + 209, \\
& 51t + 3i - 1, & 52t + 3i - 1, & 53t + 3i - 7, & 54t + 3i - 4, \\
& 55t + 3i - 1, & 56t + 3i + 746, & 59t + 4i + 157, & 59t + 4i + 215, \\
& 61t + 4i + 208, & 61t + 4i - 2, & 63t + 4i - 7, & 63t + 4i - 5, \\
& 65t + 4i - 2, & 65t + 4i + 740, & 69t + 5i + 157, & 70t + 5i + 214, \\
& 71t + 5i + 207, & 72t + 5i - 8, & 73t + 5i - 5, & 74t + 5i - 3, \\
& 75t + 5i + 739, & 80t + 6i + 156, & 80t + 6i + 213, & 82t + 6i + 201, \\
& 82t + 6i - 6, & 84t + 6i - 3, & 84t + 6i + 738, & 90t + 7i + 155, \\
& 91t + 7i + 207, & 92t + 7i + 203, & 93t + 7i - 4, & 94t + 7i + 738, \\
& 101t + 8i + 149, & 101t + 8i + 209, & 103t + 8i + 205, & 103t + 8i + 737, \\
& 111t + 9i + 151, & 112t + 9i + 211, & 113t + 9i + 946, & 122t + 10i + 153, \\
& 122t + 10i + 952, & 132t + 11i + 894.
\end{aligned}$$

B_0 is a path with an end vertex x and length $n - n_t$. Other B_i is simply a cycle of length i .

Then $f(n) \geq n + \frac{107}{3}t + \frac{7}{3}$, for $n \geq n_t$. This completes the proof.

From Theorem 7, we have

$$\liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{7654}{19071}},$$

which is better than the previous bounds $\sqrt{2}$ (see [15]), $\sqrt{2 + \frac{2}{5}}$ (see [9]).

So Conjecture 4 is not true.

Combining this with Boros, Caro, Füredi and Yuster's upper bound, namely

Theorem 2, we get

$$1.98 \geq \limsup_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \liminf_{n \rightarrow \infty} \frac{f(n) - n}{\sqrt{n}} \geq \sqrt{2 + \frac{7654}{19071}}.$$

3 Acknowledgment

The author would like to thank the referees for their many valuable comments and suggestions.

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan, New York, 1976). MR3138588
- [2] E. Boros, Y. Caro, Z. Furedi and R. Yuster, Covering non-uniform hypergraphs, J. Combin. Theory Ser. B 82(2001), 270-284. MR1842115 (2002g:05130)
- [3] G. Chen, J. Lehel, M. S. Jacobson and W. E. Shreve, Note on graphs without repeated cycle lengths, J. Graph Theory. 29(1998),11-15. MR1633908 (2000d:05059)
- [4] J. C. George, A. Khodkar, W. D. Wallis, Pancyclic and bipancyclic graphs, Springer Briefs in Mathematics, Springer, Cham, 2016. MR3495006
- [5] X. Jia, Some extremal problems on cycle distributed graphs, Congr. Numer. 121(1996), 216-222. MR1431994 (97i:05068)
- [6] C. Lai, On the size of graphs with all cycle having distinct length, Discrete Math. 122(1993) 363-364. MR1246693(94i:05048)
- [7] C. Lai, The edges in a graph in which no two cycles have the same length, J. Zhangzhou Teachers College (Natural Science) 8(4)(1994), 30-34. MR1336586 (96h:05121)
- [8] C. Lai, A lower bound for the number of edges in a graph containing no two cycles of the same length, Electron. J. Combin. 8(2001), N9, 1 - 6. MR1877662 (2002k:05124)
- [9] C. Lai, Graphs without repeated cycle lengths, Australas. J. Combin. 27(2003), 101-105. MR1955391 (2003m:05102)
- [10] C. Lai, M. Liu, Some open problems on cycles, J. Combin. Math. Combin. Comput. 91(2014), 51-64. MR3287706
- [11] S. Liu, On r -(k)-pancyclic graphs, Ars Combin. accepted.

- [12] S. C. Locke, Unsolved problems: <http://math.fau.edu/locke/Unsolved.htm>
- [13] K. Markstrom, A note on uniquely pancyclic graphs, *Australas. J. Combin.* 44 (2009), 105-110. MR2527003 (2010i:05180)
- [14] Y. Shi, Some theorems of uniquely pancyclic graphs, *Discrete Math.* 59 (1986), no. 1-2, 167-180. MR0837964
- [15] Y. Shi, On maximum cycle-distributed graphs, *Discrete Math.* 71(1988), 57-71. MR0954686 (89i:05169)
- [16] Y. Shi, The number of edges in a maximum graphs with uni-length cycles, *Kexue Tongbao.* 33(10)(1988), 795 - 796.
- [17] Y. Shi, H. P. Yap, S. K. Teo, On uniquely r -pancyclic graphs, *Graph theory and its applications: East and West* (Jinan, 1986), *Ann. New York Acad. Sci.* 576(1989), 487-499. MR1110849 (93d:05088)
- [18] Y. Shi, Some theorems on simple MCD-graphs, *J. Math. Res. Exposition* 11(4)(1991), 551-554. MR1144956 (92i:05182)
- [19] Y. Shi, The number of edges in a maximum cycle distributed graph, *Discrete Math.* 104(1992), 205-209. MR1172850 (93d:05083)
- [20] Y. Shi, On simple MCD graphs containing a subgraph homomorphic to K_4 , *Discrete Math.* 126(1994), 325-338. MR1264498 (95f:05072)
- [21] F. Tian, The progress on some problems in graph theory, *Qufu Shifan Daxue Xuebao Ziran Kexue Ban.* 1986, no. 2, 30-36. MR0865617
- [22] H. P. Yap, S. K. Teo, On uniquely r -pancyclic graphs, *Lecture Notes in Math.* 1073(1984), 334-335.
- [23] C. T. Zamfirescu, (2)-pancyclic graphs, *Discrete Appl. Math.* 161(2013), 1128-1136. MR3030597
- [24] K. Zhang, Progress of some problems in graph theory, *J. Math. Res. Exposition* 27(3) (2007), 563-576. MR2349503